

2015年12月10日

10

- 15

SVB
A22

25

- 30

35

Deh C

Sub C
4. The molecule of claim 2, in which said first random peptide library is the same library as said second random peptide library.

5 5. The molecule of claim 1, in which said method further comprises comparing the sequences of a plurality of different first peptides identified as binding said first ligand in step (a), to identify a consensus binding sequence, in which said second ligand of step (b) comprises said consensus binding
10 sequence.

Sub A23
6. The molecule of claim 2, in which said method further comprises comparing the sequences of a plurality of different first peptides identified as binding said antibody or
15 antigen-binding derivative thereof in step (a), to identify a consensus binding sequence, in which said compound of step (b) comprises said consensus binding sequence.

7. The molecule of claim 1 in which the first ligand
20 comprises a receptor.

Sub A24
8. The molecule of claim 2 in which the antibody is the monoclonal antibody 7E11-C5.

00849 01300
25 9. The molecule of claim 1 in which the library of step (a) or step (b) is a library of recombinant vectors that express a plurality of heterofunctional fusion proteins, said fusion proteins comprising a binding domain encoded by an oligonucleotide comprising unpredictable nucleotides in which
30 the unpredictable nucleotides are arranged in one or more contiguous sequences, wherein the total number of unpredictable nucleotides is greater than or equal to about 15 and less than or equal to about 600.

Sub A25
35 10. The molecule of claim 2 in which the library of step (a) or step (b) is a library of recombinant vectors that express a plurality of heterofunctional fusion proteins, said fusion proteins comprising a binding domain encoded by an oligonucleotide comprising unpredictable nucleotides in which the unpredictable nucleotides are arranged in one or more

contiguous sequences, wherein the total number of unpredictable nucleotides is greater than or equal to about 15 and less than or equal to about 600.

5 11. The molecule of claim 1 in which the library of step (a) or step (b) is a chemically synthesized library.

10 12. A molecule comprising: an amino acid sequence selected from the group consisting of:

GIINANDPLPFWFMSPTPGPAPIDINASRALVSNESE (SEQ ID NO: 1),
CGRAYCLSGNYNIFGALEPGVSTPYADVGHDDAQSWRR (SEQ ID NO: 3),
DLSRNLDGFRFLLYNAYVPGFTPTFISLTAEHLSSPKG (SEQ ID NO: 2),
RCSPIWGISYPFGLLSSNPGVCHSSDAETNIRNDILTT (SEQ ID NO: 4),

15 and

GHSNYCFVSTLGMPIVGFPSINARGLIHYGGSDPRLAA (SEQ ID NO: 5);
or a binding portion thereof.

20 13. A peptide in which the amino acid sequence of said peptide consists of the sequence selected from the group consisting of:

GIINANDPLPFWFMSPTPGPAPIDINASRALVSNESE (SEQ ID NO: 1),
CGRAYCLSGNYNIFGALEPGVSTPYADVGHDDAQSWRR (SEQ ID NO: 3),
DLSRNLDGFRFLLYNAYVPGFTPTFISLTAEHLSSPKG (SEQ ID NO: 2),
RCSPIWGISYPFGLLSSNPGVCHSSDAETNIRNDILTT (SEQ ID NO: 4),

25

and

GHSNYCFVSTLGMPIVGFPSINARGLIHYGGSDPRLAA (SEQ ID NO: 5);
or a binding portion thereof.

30

14. A method of identifying a peptide which binds to a substance of interest, comprising:

- (a) screening a first random peptide library with a ligand, said ligand being a specific binding partner of said substance of interest, to identify a first peptide that specifically binds to said ligand; and
- (b) screening a second random peptide library with a compound comprising said first peptide identified in step (a), to identify a second peptide which binds to said

35

Sub
A26

compound and which binds to said substance of interest.

15. A method of identifying a peptide which binds to
5 an antigen of interest, comprising:

- 10 (a) screening a first random peptide library with an antibody or antigen-binding derivative thereof that specifically binds to an antigen of interest, to identify a first peptide that specifically binds to said antibody or antigen-binding derivative thereof; and
- 15 (b) screening a second random peptide library with a molecule comprising said first peptide identified in step (a), to identify a second peptide sequence which binds to said molecule and which binds to said antigen of interest.

008770" 52848450
Seed
A27

20 16. The method of claim 14, in which said first random peptide library is a different library from said second random peptide library.

25 17. The method of claim 14, in which said first random peptide library is the same library as said second random peptide library.

30 18. The method of claim 14 in which the ligand is a receptor.

Sub
A28

19. The method of claim 15 in which the antibody is the monoclonal antibody 7E11-C5.

35 20. The method of claim 14 in which the library of step (a) or step (b) is a library of recombinant vectors that express a plurality of heterofunctional fusion proteins, said fusion proteins comprising a binding domain encoded by an oligonucleotide comprising unpredictable nucleotides in which the unpredictable nucleotides are arranged in one

or more contiguous sequences, wherein the total number of unpredictable nucleotides is greater than or equal to about 15 and less than or equal to about 600.

5 21. The method of claim 15 in which the library of step (a) or step (b) is a library of recombinant vectors that express a plurality of heterofunctional fusion proteins, said fusion proteins comprising a binding domain encoded by an
10 oligonucleotide comprising unpredictable nucleotides in which the unpredictable nucleotides are arranged in one or more contiguous sequences, wherein the total number of unpredictable nucleotides is greater than or equal to about 15 and less than or equal to about 600.

15 22. The method of claim 14 where the library of step (a) or step (b) is a chemically synthesized library.

20 23. A method of detecting or measuring an analyte of interest in a sample, comprising:

(a) contacting a sample with a molecule comprising a peptide capable of specifically binding said analyte of interest under conditions such that
25 specific binding between said molecule and said analyte can occur; and

(b) detecting or measuring the amount of said binding in which the presence and amount of said binding indicates the presence and amount,
30 respectively, of said analyte in the sample;

35 in which said peptide is identified by the method of claim 14.

24. The method of claim 23 in which said molecule is immobilized on a solid substratum.

008770" 62848760

Sub
A29

Sub
A30

25. A method of determining the location in a patient of a tumor comprising:

5 (a) introducing a molecule comprising a peptide that specifically binds to a tumor antigen into the patient; and

(b) determining the location in the patient of the molecule;

in which the molecule is detectably labeled; and in
10 which said peptide is identified by a method comprising:

(i) screening a first random peptide library with an antibody or antigen-binding derivative

15 thereof that
specifically binds to
said tumor antigen, to
identify a first peptide
- that specifically binds
to said antibody or
antigen-binding
derivative thereof; and

20

(ii) screening a second random peptide library with a molecule comprising said first peptide identified in (i), to identify a second peptide which binds to said molecule and which binds to said tumor antigen.

26. A therapeutic or diagnostic composition
35 comprising the molecule of claim 1; and a
pharmaceutically acceptable carrier.

27. A therapeutic or diagnostic composition comprising the molecule of claim 2; and a pharmaceutically acceptable carrier.

28. A therapeutic or diagnostic composition comprising the molecule of claim 5; and a pharmaceutically acceptable carrier.

5 29. A therapeutic or diagnostic composition comprising the molecule of claim 7; and a pharmaceutically acceptable carrier.

Sub
A32 10 30. A therapeutic or diagnostic composition comprising the molecule of claim 8; and a pharmaceutically acceptable carrier.

15 31. A therapeutic or diagnostic composition comprising the molecule of claim 12; and a pharmaceutically acceptable carrier.

20 32. A composition comprising a plurality of molecules of claim 1, in which said peptide sequences of said molecules differ.

Sub
A33 25 33. A molecule comprising a peptide or a binding portion thereof which binds to a ligand of interest, which peptide is identified by a method comprising: screening a random peptide library with a ligand of interest, said ligand of interest being a peptide having a length of between 5 and 40 amino acids, to identify a peptide that specifically binds to the ligand of interest, in which the ligand of interest is also specifically bound by an antibody or a receptor.

30 34. The molecule of claim 31 in which the ligand is a peptide having a length of between 10 and 20 amino acids.

Sub
A34 35 35. A method of obtaining an image of an internal region of a subject comprising administering to said subject an effective amount of the molecule of claim 1 in which said molecule is radiolabeled with a radioactive metal, and recording the scintigraphic image obtained from the decay of said radioactive metal.

Sub
A-34
Cont

36. A molecule comprising a peptide which binds to a substance of interest, which peptide is identified by a method comprising: screening a random peptide library with a ligand, said ligand being a peptide of 36 amino acids or fewer, in which the ligand is an epitope of an antigen that is specifically bound by an antibody or in which the ligand represents the portion of a receptor-ligand that is responsible for the specific binding of the receptor to the receptor-ligand.

37. A peptide comprising the amino acid sequence WQGTHF (SEQ ID NO: 23) and the amino acid sequence LVSKND SG (SEQ ID NO: 24) that specifically binds to an antigen of human prostate carcinoma cells.

38. A molecule comprising an amino acid sequence selected from the group consisting of:

SFMDYFRHTPEPKPAGYPNAYTDPKHPA (SEQ ID NO: 26),
SSSIFDYAPFSWGSAGLSNSSINVFERS (SEQ ID NO: 27),
SASLWDALGCWTTSAVPSYPRPHQTPGR (SEQ ID NO: 28),
SLGLPWIDVFGRSSAEPWPFGRTNLPRS (SEQ ID NO: 29),
SVHGAFLD SFFPWAADGPHGRGRLTSF (SEQ ID NO: 30),
EEKQGGRWSTMMPRPWCHEGGCGFLYYDAMTKPKTPPIMRTAA
(SEQ ID NO: 31),
LPRPFDDASWKLRAVKESPDGCGFCSPLLFPYPYGLPTFSSCD
(SEQ ID NO: 32),
GSFESARGVTCIGNHSIGAHGCGPLRSYASFNRGSGRRH (SEQ
ID NO: 33),
DQIGSRPQTTSRSISGSWENAKTLWQQDYAFSAPNAA (SEQ ID
NO: 34),
LSDAWGNFTTSYRDSAGFP SHAMTTSQGGKRNHASRFP (SEQ ID
NO: 35),
VQLDDTSPRASGQETSQSEYDARPLLSKFAIPRPWSR (SEQ ID
NO: 36),
IDSSKNRISGTGYLSFPHIRHANRRHMADDSNLAPGPS (SEQ ID
NO: 37),
WSIGTHTGPEGKFRIPCDRSGCGGTTLTHGGLNSSPTGQHERP
(SEQ ID NO: 38),
DPCEDGYWLSSVGRAGASIRGCCAIRRSSRTLTAEYSTTRASNH
(SEQ ID NO: 39),

003710 643460

GSKRSCWGTTISNYFRPVPEGCGSASSINPNTNTGRLPSLHRQ
(SEQ ID NO: 40),
SSASSGCLGRAEHLDLDSVWGCGSQADMSRRYSPWYGRPRTGV
(SEQ ID NO: 41),
5 NVMWSSSKAGIRDSCSQVPPGGCGPVNRHRASPLTPFRHGSIR
(SEQ ID NO: 42),
PLTSGSSSEYRNRDDCPVYKYATNCPRLNFSRYSPPF (SEQ ID
NO: 43),
10 GDAYGGIFSRPROGLADSYIHASYTGKHFFRGPRPPTR (SEQ ID
NO: 44),
STCIGAEGEWKSFHNFLQCRDATSTSSSTLDPTALRFG (SEQ ID
NO: 45),
YSATLWDQFGSRQVELWSNRHASSALPFASRASVLGSR (SEQ ID
NO: 46),
15 ILGWPFLTGLGDSTVHPRGRKGTDP (SEQ ID NO: 47),
SIPFSMWLNQLGSAALPSKGNQDRSD (SEQ ID NO: 48),
SRDDIFTGGPLVLFEGSKTSNHDVHSMR (SEQ ID NO: 49),
RAELVNWYEFHVTAEAEPTVINSHNMT (SEQ ID NO: 50),
GAPVWRGNPRWRGPGCFKWPGCGNGPMCNTFTPARGGSRNNGP
20 (SEQ ID NO: 51),
GSASSCFPNFTARGVTVGFFGCGSPAHPAAPRVLNPATDFPAP
(SEQ ID NO: 52),
VFRRTARSSRPIGATVFPWYGCGNSNDETLPHHDSPPSFFLGA
(SEQ ID NO: 53),
25 NTCWTDLFWHGLPGGDLPRDGCGLPSELTTHTPSRERRDASEN
(SEQ ID NO: 54),
IDWNWLERGQHNRGYLHSFPDAKSQPTRGPRVAPNGND (SEQ ID
NO: 55) and
GRGSDMREHWPWSMPLILDQHNDPSPRAQSHYYSHPF (SEQ ID
30 NO: 56).

39. A molecule comprising an amino acid
sequence selected from the group consisting of:
35 VSTGWSGTPRWCAPGGKQSSCGNGPRWTTLTPDLGGTRKYGP (SEQ ID
NO: 57),
GAPLWCEKLSGTGSGCFKWPGCGSGPTYNFTTPARVGSDNKWP (SEQ ID
NO: 58),
GPPVWSAKSRWTGTGVLNWP GCGKVRSCSTYTPSRDRSRKSDP (SEQ ID
NO: 59),

005449.04300

GSALLTSKGCVRGPGGLMRPGCGNDRLGKSSTYAHGGWIKTGP (SEQ ID
NO: 60),
GSPVWSGDNRWRGSSPLKRPGCGNGAKCNTLKDNRKDSRKTGH (SEQ ID
NO: 61),
5 G PLLPGEAAVHGARGLMRSGCGNGPTWNRLTAACRDSRNKGP (SEQ ID
NO: 62),
GSPVWMGSTRWTGHGWFRSQGCGNVPRTNSCAPAGKDSQNKGP (SEQ ID
NO: 63),
GAPVWRGNRWCSDNELERPGCGYGPRFNILPPGRGNSRKPS (SEQ ID
10 NO: 64),
GSSGWKVKHRCGGPGTLQRP GCGNPLGHTFPPTRGGSHMEGA (SEQ ID
NO: 65),
GPRSWGQPRGSDAGSCKWAGCGDAPMWRASTPGHGGPPNRGS (SEQ ID
NO: 66),
15 EALVCRGKPPWSGPAGLLWQCGTG PVSRTFTSAQGRSRNKTS (SEQ ID
NO: 67),
GAPVVG DILWCSGARGAKWPGCGKGPTNKTFSHSRGGTQKSGL (SEQ ID
NO: 68),
GAPVSRCKPACGGFWGVNWPGCGNASMCKFTNGHGVSSDNH (SEQ ID
20 NO: 69),
GAHGYKNGSTCTGLGGWRCRGCGKGAMCNPSPAGGAYHNQGP (SEQ ID
NO: 70),
G PQGSEHQCCSGHWGLKFP GCGNGPICNNFTALRGASRKNGP (SEQ ID
NO: 71),
25 GEPVWCRHSGGRVQGGLDWL GCGDGPLRYTVTPARGGPSKHGP (SEQ ID
NO: 72),
GLSLVRGDSWGSAGGWKRHGC GHPMYNPQTPARGGSCTRNT (SEQ ID
NO: 73),
VSRWSGKPRLMGSHGLNCPGCGKGHS GIMFIPDPAGSANTPP (SEQ ID
30 NO: 74),
CAPMWSGKPPWC VGGGVKFRGCGNRPDCNIITPRLVESRDKAL (SEQ ID
NO: 75) and
ADPVCSRKPDGGGLRGLRWPGCGKGPILYNVTAARGGSRNNGP (SEQ ID
35 NO: 76).

40. The molecule which binds to a ligand of interest of claim 33 in which said ligand comprises VTSAPDTRPAPGSTAPPAHGVTSAPDTR (SEQ ID NO: 9) or a portion thereof.

41. A therapeutic or diagnostic composition comprising a molecule chosen from the group of molecules of claim 38 and a pharmaceutically acceptable carrier.

5 42. A therapeutic or diagnostic composition comprising a molecule chosen from the group of molecules of claim 39 and a pharmaceutically acceptable carrier.

10 43. A molecule that binds to polymorphic epithelial mucin, comprising an amino acid sequence represented by the formula:

$R_1R_2R_3R_4R_5R_6R_7R_8R_9R_{10}R_{11}R_{12}R_{13}R_{14}R_{15}R_{16}R_{17}R_{18}R_{19}R_{20}R_{21}R_{22}R_{23}R_{24}R_{25}R_{26}$
 $R_{27}R_{28}R_{29}R_{30}R_{31}R_{32}R_{33}R_{34}R_{35}R_{36}R_{37}R_{38}R_{39}R_{40}R_{41}R_{42}R_{43}$ (SEQ ID NO: 88)

15 wherein:

$R_1 = G, C, E, \text{ or } V;$
 $R_2 = A, S, P, \text{ or } L;$
 $R_3 = P, T, H, \text{ or } L;$
20 $R_4 = L, M, Q, G, A, \text{ or } S;$
 $R_5 = W \text{ or } Y;$
 $R_6 = S, C, K \text{ or } T;$
 $R_7 = E, S, C, D, V, \text{ or } R;$
 $R_8 = N, H, K, S, \text{ or } E;$
25 $R_9 = L, H, R, N, Q, T, \text{ or } G;$
 $R_{10} = W, P, R, T, \text{ or } D;$
 $R_{11} = W, C, V, L, \text{ or } G;$
 $R_{12} = S, T, M, \text{ or } H;$
 $R_{13} = G;$
30 $R_{14} = S, A, G, N, Q, \text{ or } H;$
 $R_{15} = W, H, G, A, \text{ or } R;$
 $R_{16} = G, T, E, P, V, \text{ or } W;$
 $R_{17} = V, F, W, K, \text{ or } A;$
 $R_{18} = K, Q, D, E, R, \text{ or } L;$
35 $R_{19} = R, F, \text{ or } S;$
 $R_{20} = P, S, I \text{ or } H;$
 $R_{21} = G;$
 $R_{22} = C;$
 $R_{23} = G;$
 $R_{24} = D, S, T, N;$

- $R_{25} = G, D, L;$
 $R_{26} = P \text{ or } S;$
 $R_{27} = M, S, D, I, L, \text{ or } R;$
 $R_{28} = G, W, C, L, F, Y, \text{ or } T;$
5 $R_{29} = S, N, V, F, H, \text{ or } R;$
 $R_{30} = N, A, S, M, \text{ or } R;$
 $R_{31} = F, Q, P, \text{ or } V;$
 $R_{32} = S, V, I, K, A, \text{ or } S;$
 $R_{33} = P, A, N, \text{ or } Y;$
10 $R_{34} = G, N, \text{ or } L;$
 $R_{35} = K, R, C, Q \text{ or } L;$
 $R_{36} = V, K, R, \text{ or } A;$
 $R_{37} = G, D, A, \text{ or } E;$
 $R_{38} = S, T, P, Y \text{ or } W;$
15 $R_{39} = R, I, L, P, A \text{ or } S;$
 $R_{40} = N, K, \text{ or } M;$
 $R_{41} = S, R, T, E, Q, P, Y \text{ or } H;$
 $R_{42} = G, A, S, D, N, P, Y, \text{ or } K;$
 $R_{43} = P, H \text{ or } A. \quad -$

44. The molecule of claim 43 wherein:

- $R_1 = G;$
 $R_2 = A;$
 $R_3 = P;$
25 $R_5 = W;$
 $R_6 = S;$
 $R_{10} = W;$
 $R_{11} = W;$
 $R_{12} = S \text{ or } T;$
30 $R_{14} = S;$
 $R_{16} = G;$
 $R_{18} = K;$
 $R_{19} = R;$
 $R_{20} = P;$
35 $R_{26} = P;$
 $R_{28} = G \text{ or } W;$
 $R_{30} = N;$
 $R_{31} = F;$
 $R_{33} = P;$
 $R_{35} = K \text{ or } R;$

R₃₇ = G;
R₃₈ = S;
R₄₀ = N or K;
R₄₁ = G;

5

45. The molecule of claim 2 in which the antibody or antigen-binding derivative thereof is capable of specifically binding to a human tumor antigen.

10

Add
A35

15

20

25

30

35

00870" 62848460